2075 Bhadra

Exam.	R	egular	
Level	BE	Full Marks	80
Programme	BEL,BEX, BCT, BGE	Pass Marks	32
Year / Part	II / II	Time	3 hrs.

Subject: - Applied Mathematics (SH551)

- \checkmark Candidates are required to give their answers in their own words as far as practicable.
- ✓ Attempt <u>All</u> questions.
- ✓ The figures in the margin indicate Full Marks.
- ✓ Assume suitable data if necessary.
- 1. Define harmonic function of complex variable. Show that $u(x, y) = y^3 3x^2y$ is harmonic [1+4]and find corresponding analytic function. Define conformal mapping for function of complex variable. Show that function of complex variable w = iz is transformed through an angle $\frac{\pi}{2}$ in w-plane. [1+4] [5] 3. State and prove Cauchy's integral theorem. 4. Define Laurent's Series for the function of complex variable. Find Laurent's series of the function $f(z) = \frac{z}{(z+2)(z+3)}$ in the region 2 < |z| < 3. [1+4] 5. Define pole of order m for function of complex variable. Find residues of $f(z) = \frac{-z^2 - 2z}{(z+1)^2(z^2+1)}$ at its poles. [1+4]6. Evaluate $\int_{-\infty}^{\infty} \frac{x^2}{(x^2+1)(x^2+4)} dx$ by contour integration in the complex plane. [5] [3+2]7. Find the Z-transform of: i) $t^2 e^{at}$ ii) e^{-at} cos wt [2.5+2.5] 8. Find the inverse Z-transform of: i) $X(z) = \frac{2z^2 - 5z}{(z-2)(z-3)}$ (By partial fraction method) ii) $X(z) = \frac{z^{-2}}{(1-z^{-1})^3}$ (By inversion integral method) 9. State final value theorem for Z-transform. Obtain Z-transform of $(1 - e^{-at})$; a>0 and hence [1+4] evaluate $x(\infty)$ by using final value theorem. [5] 10. Solve the difference equation: x(k+2) - 3x(k+1) + 2x(k) = 0; given that x(0) = 0 and x(1) = 1 by using z-transform method. [5] 11. Find the Fourier integral of the function:
 - $f(x) = \begin{cases} 1, & \text{for } 0 < x < \pi \\ 0, & \text{for } x > \pi \end{cases}$

- 12. Find the Fourier transform of e^{-x^2} . Also verify the convolution theorem for $f(x) = e^{-x^2}$ and $g(x) = e^{-x^2}$ [5]
- 13. Derive one dimensional wave equation and solve it completely. [10]

14. Solve completely the Laplace equation
$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$
 under the conditions: [10]

$$u(0,y) = u(1,y) = u(x,o) = 0, u(x,\infty) = sin\left(\frac{n\pi x}{l}\right)$$

14	TRIBHUVAN UNIVERSITY
INS	TITUTE OF ENGINEERING
Exam	ination Control Division

Exam.	Back			
Level	BE	Full Marks	80	
Programme	BGE, BEL, BEX, BCT	Pass Marks	32	
Year / Part	П/П	Time	3 hrs.	

2075 Baisakh

Subject: -	- Applied	Mathematics	(SH551)	1
------------	-----------	-------------	---------	---

- ✓ Candidates are required to give their answers in their own words as far as practicable.
- ✓ Attempt <u>All questions</u>.
- The figures in the margin indicate <u>Full Marks</u>.
- ✓ Assume suitable data if necessary.
- 1. a) Define harmonic function. Is V = arg(z) is harmonic? If yes, find a corresponding harmonic conjugate. [1+1+3]
 - b) Define conformal mapping. Find the bilinear transformation which maps the points $z = 0, 1, \infty$ into the points w = -3, -1, 1 respectively. [1+4]

2. a) Distinguish between Cauchy integral Theorem and Cauchy integral formula. Using

Cauchy integral formula evaluate
$$\int_{c} \frac{e^{-z}}{(z+1)(z-2)} dz$$
 where C is the circle $|z-1|=3$. [1+4]

b) State and Prove Taylor's series for function of complex variable.

3. a) Define an isolated pole. Using Cauchy's residue theorem evaluate $\int_{c} \frac{z-1}{(z+1)^2(z-2)} dz$ where C is the circle |z-i|=2.

$$\int_{-\infty}^{\infty} \frac{x^2}{(1+x^2)(x^2+4)} dx$$

- 4. a) Obtain the z-transform of (1-e^{-at}), a > 0 and hence evaluate x(∞) by using final value theorem. [2+3]
 - b) Obtain the inverse z-transform of:

$$X(z) = \frac{2z^3 + z}{(z-2)^2(z-1)}$$
 by using partial fraction method. [5]

5. a) Define z-transform of function f(t). Find the z-transform of following sequences: [1+2+2]

- (i) $f(k) = \begin{cases} 15,10,7,4,1,-1,3,6 \\ \uparrow \end{cases}$ (ii) $f(k) = \begin{cases} 5^{k} ; k < 0 \\ 2^{k} ; k \ge 0 \end{cases}$
- b) Solve the difference equation by the application of z-transform: x(k+2)+3x(k+1)+2x(k)=0 with conditions x(0) = 0, x(1) = 1.

[5]

[5]

[5]

- 6. a) A tightly stretched string with fixed ends at x = 0 and x = 1 is initially at rest in its equilibrium position. Find the deflection u(x, t) if it is set vibrating by giving to each of its points a velocity $3(lx-x^2)$.
 - b) Derive two dimensional heat equation.
- 7. a) Obtain the Fourier sine integral representation of $e^{-x}\cos x$ and hence show that $\int_{0}^{\infty} \frac{\omega^{3} \sin \omega x}{\omega^{4} + 4} d\omega = \frac{\pi}{2} e^{-x} \cos x, \quad x > 0.$
 - b) Find the Fourier Cosine transform of $f(x) = e^{-x}$, x > 0 and hence by Parseval's identity, show that

$$\int_0^\infty \frac{1}{(1+x^2)^2} dx = \frac{\pi}{4} \; .$$

[5]

[10]

[10]

Exam.		Regular	
Level	BE	Full Marks	80
Programme	BGE, BEL, BEX, BCT	Pass Marks	32
Year / Part	II / II	Time	3 hrs.

2074 Bhadra

Subject: - Applied Mathematics (SH551)

- ✓ Candidates are required to give their answers in their own words as far as practicable.
- ✓ Attempt <u>All</u> questions.
- ✓ The figures in the margin indicate <u>Full Marks</u>.
- ✓ Assume suitable data if necessary.
- a) Define an analytic function for a function of complex variable. Derive Cauchy Riemann equations in Cartesian form. [1+4]
 b) Define linear fractional mapping. Find bilinear mapping which maps the points z = 0, 1, -1 to w = i, 2, 4. [1+4]
 a) State and Prove Cauchy integral theorem. [5]
 b) Point out difference between Taylor's series and Laurent's series. Find Laurent' series [5]

of function
$$f(z) = \frac{\sin z}{z^6}$$
, $0 < |z| < TR$ [1+4]

- 3. a) Define pole of order m. Using Cauchy's residue theorem evaluate $\int \cot z \, dz$; where C is |z| = 1. [1+4]
 - b) Using Counter integration evaluate,

$$\int_{-\infty}^{\infty} \frac{\mathrm{d}x}{\left(1+x^2\right)^2} \,. \tag{5}$$

4. a) Find the z-transform of:

(i) cosat (ii) te^{-at} [2+3]

b) State final value theorem. If x(t) = 0 for t < 0 and Z[x(t)] = X(z) for $t \ge 0$ then prove that:

$$Z[\mathbf{x}(\mathbf{t}+\mathbf{n}\mathbf{T})] = \mathbf{z}^{\mathbf{n}} \left[\mathbf{X}(\mathbf{z}) - \sum_{k=0}^{\mathbf{n}-1} \mathbf{x}(\mathbf{k}\mathbf{T})\mathbf{z}^{k} \right].$$
[1+4]

5. a) Obtain inverse Z-transform of
$$\frac{z(3z^2 - 6z + 4)}{(z-1)^2(z-2)}$$
. [5]

- b) Solve the difference equation by the application of z-transform: x (k+2) -4x (k+1) + 4x(k) = 0; with conditions x(0) = 1; x(1) = 0. [5]
- 6. a) Derive one dimensional wave equation and solve it completely. [5+5]
 - b) A uniform rod of length ℓ has its end maintained at a temperature 0°C and the initial temperature of the rod is:

$$u(x,0) = 3\sin\frac{\pi x}{\ell} \quad \text{for } 0 < x < \ell \,.$$

Find the temperature u(x, t).

7. a) Find Fourier integral of the function

$$f(x) = \begin{cases} 1 & \text{if } |x| < 1 \\ 0 & \text{if } |x| > 1 \end{cases}$$

b) Verify the convolution theorem for Fourier transform for the functions

[10]

[5]

2073 Bhadra

Exam.	Regular			
Level	BE	Full Marks	80	
Programme	BEL, BEX, BCT, BGE	Pass Marks	32	
Year / Part	II / II	Time	3 hrs.	

[5]

[1+1+1.5+1.5]

[5]

Subject: - Applied Mathematics (SH551)

- ✓ Candidates are required to give their answers in their own words as far as practicable.
- ✓ Attempt <u>All</u> questions.
- ✓ The figures in the margin indicate *Full Marks*.
- ✓ Assume suitable data if necessary.

1. a) Define harmonic function of complex variable. Determine the analytical function

$$f(z) = u + iv \text{ if } u = y^3 - 3x^2y$$
 [1+4]

- b) Derive Cauchy-Reimann equations if function of complex variable f(z) = u + iv is analytic in cartesian form.
 [5]
- a) What do you mean by conformal mapping? Find the linear transformation which maps points z₁ = 1, z₂ = i, z₃ = -1 into the points w₁ = 0, w₂ = 1, w₃ = ∞. [1+4]
 - b) State and prove Cauchy's integral formula.
- 3. a) State Taylor's theorem. Find the Laurent's series representation of the function

$$f(z) = \frac{z}{(z+1)(z+2)}$$
 in the annular region between $|z| = 1$ and $|z| = 2$. [1+4]

b) Define zero of order m of function of complex variable .Determine the poles and residue at poles of the functions $f(z) = \frac{1+z}{(z+2)(1-z)^2}$. [1+4]

OR

Evaluate the real integral $\int_{-\infty}^{\infty} \frac{x^2}{(1+x^2)^3} dx$ by contour integration in the complex plane. [5]

4. a) Define z-transform. How does it differ from Fourier transform? Obtain z-transform of

- (i) $t^2 a^t$ (ii) cosat
- b) State initial value theorem for z transform. Find the initial value x(0) and x(1) for the function.
 [1+4]

$$X(z) = \frac{(1 - e^{T})z^{-1}}{(1 - z^{-1})(1 - e^{-T}z^{-1})}$$

- 5. a) Obtain the inverse z-transform of $X(z) = \frac{3z^3 + 2z}{(z-3)^2(z-2)}$ by using inversion integral method.
 - b) Apply method of z-transform to solve the difference equation [5] $x(k+2)^{\frac{1}{2}}-4x(k+1)+4x(k)=0; x(0)=0, x(1)=1$

6. Solve completely one-dimensional wave equation $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$ under the conditions: [10]

$$u(0,t) = 0, u(l,t) = 0, u(x,0) = 0 \text{ and } \left(\frac{\partial u}{\partial t}\right)_{at t=0} = 3(lx - x^2)$$

7. Derive one dimensional heat equation and solve it completely. [10]

- 8. a) State convolution theorem for Fourier transform. Give its importance with suitable [2+3]
 - b) Find the Fourier cosine integral of the function $f(x) = e^{-kx} (x > 0, k > 0)$ and hence show that $\int_0^\infty \frac{\cos \omega x d\omega}{k^2 + \omega^2} = \frac{\pi}{2k} e^{-kx}; x > 0, k > 0$ [5]

15 TRIBHUVAN UNIVERSITY INSTITUTE OF ENGINEERING

Examination Control Division

2072 Ashwin

Exam.	Ro	gular	
Level	BE	Full Marks	80
Programme	BEL, BEX, BCT, BGE	Pass Marks	32
Year / Part	II / II	Time	3 hrs.

[5]

[5]

[5]

[10]

[10]

[5]

Subject: - Applied Mathematics (SH551)

- ✓ Candidates are required to give their answers in their own words as far as practicable.
- ✓ Attempt <u>All</u> questions.
- The figures in the margin indicate <u>Full Marks</u>.
- Assume suitable data if necessary.
- 1. a) If $u = (x-1)^3 3xy^2 + 3y^2$, determine V so that u + iv is an analytic function of x+iy. [5]
 - b) Define an analytic function. Express Cauchy Riemann equations $u_x = v_y$ and $u_y = -v_x$ in polar from.
- 2. a) Find the bilinear transformation which maps points z₁ = 1, z₂ = i, z₃ = -1 into the points w₁ = i, w₂ = -1, w₃ = -i respectively. [5]
 - b) Evaluate $\int_{0}^{1+i} (x^2 + iy) dz$ along the path $y = x^2$ [5]

3. a) Express
$$f(z) = \frac{1}{(z^2 - 3z + 2)}$$
 as Laurent's series in the region $1 < |z| < 2$. [5]

b) Evalute
$$\int_{0}^{2\pi} \frac{1}{5-4\sin\theta} d\theta$$
 by contour integration method in complex plane. [5]

- 4. a) Find z-transform of:
 - i) te^{-at}
 - ii) sinat
 - b) State and prove final value theorem for z- transform.

5. a) Find the inverse z-transform of $\frac{2z^2 - 5z}{(z-2)(z-3)}$ by using partial fraction method. [5]

- b) Solve difference equation $x(k+2)-3x(k+1)+2x(k) = 4^k$ for x(0) = 0 and x(1) = 1. [5]
- 6. Derive one dimensional wave equation and obtain its solution.
- 7. Solve one dimensional heat equation:

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$$
 under the conditions:

i) u is not infinite as $t \rightarrow \infty$

- *ii)* $\frac{\partial u}{\partial x} = 0$ for x = 0 and x = l
- iii) $u(x,0) = lx x^2$ for t = 0; between x = 0 and x = l

8. a) Find Fourier integral representation of $f(x) = e^{-x}, x > 0$ and hence evaluate $\int_{0}^{\infty} \frac{\cos(sx)}{s^{2} + 1} ds$ [5]

b) Find the Fourier cosine transform of $f(x) = e^{-|x|}$ and hence, by Parseval's identity, shown that $\int_0^{\infty} \frac{1}{(1+x^2)^2} dx = \frac{\pi}{4}$

*** 9

2071 Bhadra

Exam.	Regular / Back		
Level	BE	Full Marks	80
Programme	BEL, BEX, BCT, BGE	Pass Marks	32
Year / Part	П/П	Time	3 hrs.

Subject: - Applied Mathematics (SH551)

- \checkmark Candidates are required to give their answers in their own words as far as practicable.
- ✓ Attempt <u>All</u> questions.
- ✓ <u>All</u> questions carry equal marks.
- ✓ Assume suitable data if necessary.
- 1. Determine the analytic function f(z) = u + iv if $u = \log \sqrt{x^2 + y^2}$.
- 2. State and prove Cauchy's integral formula.
- 3. Find the Taylor's series of $f(z) = \frac{1}{1-z}$ about z = 3i.
- 4. Evaluate the integral: $\oint_C \frac{z^2 dz}{(z+1)(z+3)}$ where C: |z| = 4, using residue theorem.
- 5. Define conformal mapping, show that $w = \frac{az+b}{cz+d}$ is invariant to

$$\left(\frac{\mathbf{w}-\mathbf{w}_1}{\mathbf{w}-\mathbf{w}_3}\right) \times \left(\frac{\mathbf{w}_2-\mathbf{w}_3}{\mathbf{w}_2-\mathbf{w}_1}\right) = \left(\frac{\mathbf{z}-\mathbf{z}_1}{\mathbf{z}-\mathbf{z}_3}\right) \times \left(\frac{\mathbf{z}_2-\mathbf{z}_3}{\mathbf{z}_2-\mathbf{z}_1}\right)$$

6. Using contour integration, evaluate real integral:

$$\int_{-\infty}^{+\infty} \frac{x^2 dx}{(x^2 + a^2)(x^2 + b^2)}$$

7. Find the z-transform of $x(z) = \cosh t \sinh t$.

8. State and prove "final value theorem" for the z-transform.

- 9. Find the inverse z-transform of $x(z) = \frac{z}{z^2 + 7z + 10}$.
- 10. Using z-transform solve the difference equation:

$$x(K+2) + 6x(K+1) + 9x(K) = 2^{K}; x_0 = x_1 = 0.$$

- 11. Derive one-dimensional heat equation.
- 12. Solve the wave equation for a tightly stretched string of length 'l' fixed at both ends if the initial deflection in $y(x, 0) = |x x^2|$ and the initial velocity is zero.

13. Solve
$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{2y^2} = 0$$
 under the conditions $u(0, y) = u(1, y) = u(x, 0) = 0$, $u(x, a) = \sin\left(\frac{n\pi x}{1}\right)$

- 14. Derive the wave equation (vibrating of a string).
- 15. Find the Fourier cosine transform of $f(x) = e^{-|m|x}$ and hence show that $\int_{0}^{\infty} \frac{\cos p\gamma}{\gamma^2 + \beta^2} d\gamma = \frac{\pi}{2\beta} e^{-p\beta}$.

16. Find the Fourier integral representation of the function $f(x) = e^{-x}$, $x \ge 0$ with f(-x) = f(x). Hence evaluate $\int_{0}^{\infty} \frac{\cos(sx)}{s^{2}+1} ds$.



|13 N

12 TRIBHUVAN UNIVERSITY INSTITUTE OF ENGINEERING

Examination Control Division

2071 Magh

Exam.	New Back (2066 & Later Batc				
Level	BE	Full Marks	80		
Programme	BEL, BEX, BCT, BGE	Pass Marks	32		
Year / Part	II / II	Time	3 hrs.		

Subject: - Applied Mathematics (SH551)

- ✓ Candidates are required to give their answers in their own words as far as practicable.
- ✓ Attempt <u>All</u> questions.

ANG STORAGE

- ✓ The figures in the margin indicate <u>Full Marks</u>.
- ✓ Assume suitable data if necessary.

1. a) Determine the analytic function
$$f(z) = u + iv$$
 if $u = 3x^2y - y^3$. [5]

b) Find the linear transformation which maps the points $z = 0, 1, \infty$ into the points w = -3, -1, 1 respectively. Find also fixed points of the transformation. [5]

2. a) State and prove Cauchy's integral formula.

b) Evaluate
$$\int_{C} \frac{e^{2z}}{(z-1)(z-2)} dz$$
 where C is the circle $|z| = 3$. [5]

3. a) Find the first four terms of the Taylor's series expansion of the complex function
$$f(z) = \frac{z+1}{(z-3)(z-4)}$$
 about the centre z = 2. [5]

b) Evaluate
$$\int_{C} \frac{4-3z}{z(z-1)(z-2)} dz$$
 where C is the circle $|z| = \frac{3}{2}$. [5]

OR

Evaluate $\int_0^{2\pi} \frac{1}{\cos \theta + 2} d\theta$ by contour integration in the complex plane.

4. Derive one dimensional heat equation $u_t = c^2 u_{xx}$ and solve it completely. [10]

- 5. Find all possible solution of Laplace equation $u_{xx} + u_{yy} = 0$. Using this, hence solve $u_{xx} + u_{yy} = 0$, under the conditions u(0, y) = 0, u(x, y) = 0 when $y \to \infty$ and $u(x, 0) = \sin x$. [10]
- 6. a) Find the z-transform of sin K θ . Use it to find the $z[a^{K} \sin K\theta]$. [5]

b) If
$$z[x(K)] = \frac{2z^2 + 3z + 12}{(z-1)^4}$$
, find the value of x(2) and x(3). [5]

7. a) Find the inverse z-transform of $x(z) = \frac{3z^3 + 2z}{(z-3)^2(z-2)}$ by using inversion integral method.

b) Using z-transform solve the difference equation $x(K + 2) - 4x(K + 1) + 4x(K) = 2^{K}$ given that x(0) = 0, x(1) = 1. [5]

[5]

[5]

8. a) Find the Fourier sine integral of the function $f(x) = e^{-Kx}$ and hence show that $\int_{0}^{\infty} \lambda \sin \lambda x$ $\pi = \frac{\pi}{K} x$

$$\int_{0}^{\frac{\lambda \sin \lambda x}{\lambda^{2} + \beta^{2}}} d\lambda = \frac{\pi}{2} e^{-Kx}, \quad x > 0, K > 0$$

b) Find the Fourier sine transform of
$$e^{-x}$$
, $x \ge 0$ and hence show that [5]

$$\int_{-\infty}^{\infty} \frac{x \sin mx}{x^2 + 1} dx = \frac{\pi}{2} e^{-m}, m \ge 0$$

25 TRIBHUVAN UNIVERSITY	Exam.		Regular	
INSTITUTE OF ENGINEERING	Level	BE	Full Marks	80
Examination Control Division	Programme	BEL, BEX, BCT	Pass Marks	32
2070 Bhadra	Year / Part	II / II	Time	3 hrs.

Subject: - Applied Mathematics (SH551)

 \checkmark Candidates are required to give their answers in their own words as far as practicable.

✓ Attempt All questions.

✓ The figures in the margin indicate <u>Full Marks</u>.

✓ Assume suitable data if necessary.

27

- Show that u(x, y) = x² + 2xy y² is a harmonic function and determine v(x, y) in such a way that f(z) = u(x, y) + iv(x, y) is analytic.
- 2. Define complex integral. State and prove Cauchy integral formula.

OR

[5]

[5]

[5]

Obtain bilinear transformation which maps -i, o, i to -1, i, 1.

3. Evaluate
$$\int_C \frac{e^{zz}}{(z-1)(z-2)} dz$$
 where C is $|z| = 3$ using Cauchy's integral formula. [5]

4. Obtain the Laurent series which represents the function $f(z) = \frac{z^2 - 1}{(z+2)(z+3)}$ 2 < |z| < 3. [5]

5. Find the Laurent series of
$$f(z) = \frac{1}{4+z^2}$$
 about the point $z = i$. [5]

- 6. State and prove Taylor series of a function f(z).
- 7. Derive one dimensional wave equation $u_{tt} = c^2 u_{xx}$ and solve it completely. [10]
- 8. Solve one dimensional heat equation $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$ under the boundary condition $\frac{\partial u}{\partial x} = 0$ when x = 0 and x = L and initial condition u(x, 0) = x for 0 < x < L. [10]

9. Find Z transform of (a)
$$te^{-at}$$
 and (b) sin at. [5]

10. Find the inverse z-transform (a)
$$\frac{z-4}{(z-1)(z-2)^2}$$
 (b) $\frac{z}{z^2-3z+2}$. [5]

- 11. Obtain the Z transform of $x(t) = (1 e^{-at})$, a > 0 and hence evaluate $x(\infty)$ by using final value theorem. [5]
- 12. Solve using z-transform the difference equation x(K + 2) + 2x(K + 1) + 3x(K) = 0. [5]
- 13. Find the Fourier sine transform of $f(x) = e^{-x}$, $x \ge 0$ and hence evaluate $\int_0^\infty \frac{x \sin x}{(1+x^2)} dx$. [5]
- 14. State and prove convolution theorem of Fourier transform.

[5]

25 TRIBHUVAN UNIVERSITY Exam. New Back (2066 & Later Batch) INSTITUTE OF ENGINEERING Level BE Full Marks BEL, BEX, Examination Control Division Programme **Pass Marks** BCT

Year / Part 2070 Magh II / II Time 3 hrs. Subject: - Applied Mathematics (SH551) \checkmark Candidates are required to give their answers in their own words as far as practicable. ✓ Attempt All questions. ✓ The figures in the margin indicate Full Marks. ✓ Assume suitable data if necessary. 1. Define analytic function. Show that the function $f(z) = \frac{1}{z^4}$ is analytic except z = 0[5] 2. Define complex integral. Evaluate $\int_{c} \log z \, dz; c: |z| = 1$ [5] Obtain a bilinear transformation which maps -i, 0, i to -1, i, 1. 3. Evaluate $\int_0^{1+i} (x^2 + iy) dz$ along the path y = x. [5] 4. Find the Taylor series of $f(z) = \frac{1}{4+z^2}$ about the point z = i. [5]

- 5. Evaluate the integrals by residue theorem $\int_{c} \frac{1-\cos z}{z^{3}} dz$
- 6. State Cauchy's Residue theorem and use it to evaluate $\int_{c} \frac{z^2}{3+4z+z^2} dz$ where C is |z|=2[5]

OR Evaluate $\int_{0}^{2\pi} \frac{d\theta}{\cos \theta + 2}$ by contour integration in complex plane.

- 7. Derive the one dimensional wave equation.
- 8. A rod of length L has its ends A and B maintained at 0° and 100° respectively until steady state prevails. If the changes are made by reducing the temperature of end B to 85° and increasing that of end A to 15°, then find the temperature distribution in the rod at a time t.
- 9. Find the z-transform of (i) e^{-at} sinwt (ii) cos at

10. Obtain inverse Z-transform of (i)
$$\frac{z+2}{(z-2)(z-3)}$$
, (ii) $\frac{z}{(z-2)(z-1)}$ [5]

11. If x(k) = 0 for k < 0 and $Z\{x(k)\} = X(z)$ for k > 0 then prove that $Z\{x(k+n)\} = z^n X(z) - z^n$ $\sum_{k=0}^{n-1} \chi(k) z^{-k}$ where n = 0, 1, 2.... [5]

12. Solve the difference equation x (k+2) - 4x(k+1) + 4x(k) = 0 with conditions, x(0) = 0, x(1) = 1

13. Find the cosine transform of $f(x) = e^{-mx} m > 0$ show that $\int_0^\infty \frac{\cos pr}{r^2 + B^2} = \frac{\Pi}{2B} e^{-PB}$ [5]

14. Find the Fourier transform of
$$g(x) = \begin{cases} 1 - x^2 \\ 0, \end{cases}$$
 if $-1 < x < 1;$ [5]

if otherwise. and hence use it to evaluate $\int_0^\infty \left(\frac{x\cos x - \sin x}{x^3}\right)\cos(x/2)dx$ ***

[10]

[5]

80

32

[5]

[5]

[10]

25 TRIBHUVAN UNIVERSITY	Exam.	Regular (20	66 & Later B	atch)
INSTITUTE OF ENGINEERING	Level	BE	Full Marks	80
Examination Control Division	Programme	BEL, BEX, BCT	Pass Marks	32
2069 Bhadra	Year / Part	II / II	Time	3 hrs.

Subject: - Applied Mathematics (SH551)

Candidates are required to give their answers in their own words as far as practicable.

✓ Attempt <u>All</u> questions.

 \checkmark <u>All</u> questions carry equal marks.

✓ Assume suitable data if necessary.

1. Determine the analytic function f(z) = u(x,y) + iv(x,y) if $u(x,y) = x^2 - y^2$.

2. Define complex integral. Evaluate: $\oint_C (z+1)dz$ where C is the square with vertices at z = c

0, z = 1, z = 1+i and z = i.

OR

Find linear fractional transformation mapping of: $-2 \mapsto \infty, 0 \mapsto \frac{1}{2}, 2 \mapsto \frac{3}{4}$.

3. a) State Cauchy's integral formula and evaluate the integral $\oint \frac{4-3z}{z(z-1)(z-2)} dz$, where C is circle $|z| = \frac{3}{2}$.

b) Obtain the Laurent series which represents the function $f(z) = \frac{1}{(1+z^2)(z+2)}$ when $|\langle |z| < 2$.

4. a) Find the Taylor's series expansion of $f(z) = \frac{1}{z^2 + 4}$ about the point z = i.

b) Evaluate $\int \tan z \, dz$ where C is a circle |z| = 2 by Cauchy's residue theorem.

OR

Evaluate $\int_0^{2\pi} \frac{\cos 3\theta}{5 - 4\cos \theta} d\theta$ by contour integration in the complex plane.

5. Find the z-transforms of: (i) $\cos h(at) \sin (bt)$ (ii) n.(n-1); n = k

6. Find the inverse z-transforms of: (i) $\frac{Z}{Z^2 - 3Z + 2}$ (ii) $\frac{Z}{(Z+1)^2(Z-1)}$.

7. a) State and prove convolution theorem for z-transform.

b) Solve by using z-transform the difference equation x(k+2) + 2x(k+1) + 3x(k) = 0, given that x(0) = 0 and x(1) = 2

- 8. Solve $\frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$ given that u = 0 as $t \to \infty$ as well as u = 0 at x = 0 and x = l.
- 9. Solve $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$, which satisfies the condition u(0,y) = u(L,y) = u(x,0) = 0 and $u(x,a) = \sin\left(\frac{n\pi x}{L}\right)$.

OR

The diameter of a semi-circular plate of radius a is kept at 0°C and the temperature at the semi-circular boundary is u_0 Find the steady state temperature in the plate.

10. Find the Fourier integral representation of the function $f(x) = e^{-x}$, $x \ge 0$ with f(-x) = f(x).

2

Hence evaluate
$$\int_{0}^{\infty} \frac{\cos(sx)}{s^2 + 1} ds$$
.

- 11. Find the Fourier transform of: $f(x) = 1-x^2$, |x| < 1
 - = 0, $|\mathbf{x}| > 1$ and hence evaluate

$$\int_{0}^{\infty} \left(\frac{x \cos x - \sin x}{x^3} \right) \cos \frac{x}{2} dx.$$

		Pa			,	
25 TRIBHUV	AN UNIVERSITY	Exam.		Regular		
INSTITUTE OF		Level	BEL BEX	Full Marks	80	
Examination	ontrol Division	Programme.	BCT	Pass Marks	32	
2068 1	Bhadra	Year / Part	П/П	Time	3 hrs.	
	Subiect: - A	pplied Mather	natics			
. Condidatas ara r	aquirad to give their or		da oo fo		rang na tao ming na Kaoming na K	
✓ Attempt All ques	equited to give their a	iswers in their ov	vn words as fa	r as practicable	•	
✓ <u>All</u> questions can	rry equal marks.					
✓ Assume suitable	data if necessary.					
1			1 () (1		<i>.</i> •	
$f(z) = \log z$ i	ary conditions for a fission of a	unction $f(z)$ to be except at the original	e analytic. Sho	ow that the fun	iction	
h) Find the line	ar fractional transform	ation that man	the noints 7 -	$-i$ $z_{e} = 0$ and	$7_{2} = i$	
into points w	$w_1 = -1, w_2 = i, w_3 = 1$	respectively.	the points $z_1 -$	-1, 22 - 0 and	23 - 1	
$2 \sim a$) State and pro	ove Cauchy's integral	formula				
b) Write the st	atement of Cauchy's	integral formul	a lice it to e	valuate the int	teoral	
e^{z}	atoment of Cauchy s	integral formul		valuate the m	iograf	
$d\frac{z}{(z-1)(z-1)}$	dz where C is the ci	rcle $ z = 2$.				
3 a) Write the st	atement of Taylor's	theorem Find th	o Touront cor	ion for the fur	oction	
3. a) while the st	atchieft of Taylor S	meorem. rind u	ie Laureni sen			
$f(z) = \frac{1}{z^2 - 3z}$	$\frac{1}{z+2}$ in the region $1 < \frac{1}{z+2}$	z < 2.				
		<u>م</u>	inz,	<u> </u>	. •	
b) State Cauchy	-residue theorem. Usi	ng it evaluate g-	$\frac{1}{z^6}$ dz where	C: z = 1.		
		OR	•			
				ar a constant	i de la constance en la constan La constance en la constance en	
2π.	A TA CALL MARK			•		
Evaluate $\int_{-2\pi}^{2\pi}$	$\frac{d\theta}{d\theta}$ d θ by contour	integration in the	e complex plan	C.		
Evaluate $\int_{0}^{2\pi} \frac{2\pi}{2}$	$\frac{d\theta}{d\theta}$ d θ by contour $+\cos\theta$	integration in the	complex plan	с.		
Evaluate $\int_{0}^{2\pi} \frac{1}{2}$ 4. a) Show that t	$\frac{d\theta}{d\theta} = d\theta \text{ by contour}$ + cos θ he Z-transform of co	integration in the os k θ is $\frac{z(z-z)}{z}$	$\frac{1}{\cos\theta}$. Use	e. e this result to	o find	
Evaluate $\int_{0}^{2\pi} \frac{1}{2}$ 4. a) Show that t	$\frac{d\theta}{d\theta}$ d θ by contour + cos θ he Z-transform of co	integration in the os k θ is $\frac{z(z-z)}{z^2-2z}$	$\frac{\cos\theta}{\cos\theta+1}$. Use	e.	find	
Evaluate $\int_{0}^{2\pi} \frac{1}{2}$ 4. a) Show that t Z-transform	$\frac{d\theta}{d\theta} = d\theta \text{ by contour}$ + cos θ he Z-transform of co of $a^k \cos k\theta$.	integration in the os k θ is $\frac{z(z-z)}{z^2-2z}$	$\frac{\cos\theta}{\cos\theta+1}$. Use	e. e this result to	o find	
Evaluate $\int_{0}^{2\pi} \frac{1}{2}$ 4. a) Show that t Z-transform b) Obtain the ir	$\frac{d\theta}{d\theta} = d\theta \text{ by contour}$ + cos θ he Z-transform of co of $a^k \cos k\theta$.	integration in the ps k θ is $\frac{z(z-z)}{z^2-2z}$ $\frac{2z^3+z}{z^2-2z}$	$\frac{\cos\theta}{\cos\theta+1}$. Use $\sin \theta$ partial fra	e this result to action method.`	o find	
Evaluate $\int_{0}^{2\pi} \frac{1}{2}$ 4. a) Show that t Z-transform b) Obtain the in	$\frac{d\theta}{d\theta} = d\theta \text{ by contour}$ $\frac{d\theta}{d\theta} = d\theta \text{ by contour}$ he Z-transform of co of a ^k cos k θ . averse Z-transform of	integration in the ps k θ is $\frac{z(z-z)^2}{z^2-2z}$ $\frac{2z^3+z}{(z-2)^2(z-1)}$, u	$\frac{\cos\theta}{\cos\theta+1}$. Use $\sin \theta$ partial fra	e. this result to action method.`) find	
Evaluate $\int_{0}^{2\pi} \frac{1}{2}$ 4. a) Show that t Z-transform b) Obtain the in 5. a) Solve the difference	$\frac{d\theta}{d\theta} = d\theta \text{ by contour}$	integration in the ps k θ is $\frac{z(z-z)}{z^2-2z}$ $\frac{2z^3+z}{(z-2)^2(z-1)}$, w +2) - x(k+1)	$\frac{\cos \theta}{\cos \theta + 1}$. Use $\sin \theta$ partial fra + 0.25x(k) =	e. e this result to action method.` u(k) where x((() find $(0) = 1$	
Evaluate $\int_{0}^{2\pi} \frac{1}{2}$ 4. a) Show that t Z-transform b) Obtain the in 5. a) Solve the dia and $x(1) = 2$	$\frac{d\theta}{d\theta} = d\theta \text{ by contour}$ he Z-transform of co of a ^k cos k θ . averse Z-transform of fference equation x(k and u(k) is unit step fi	integration in the ps k θ is $\frac{z(z-z)}{z^2-2z}$ $\frac{2z^3+z}{(z-2)^2(z-1)}$, where $z = -2z^3$ $+2z^3+z$	$\frac{\cos\theta}{\cos\theta+1}$. Use sing partial fra + 0.25x(k) =	e. this result to action method.` u(k) where x(0	find)) = 1	· •
Evaluate $\int_{0}^{2\pi} \frac{1}{2}$ 4. a) Show that t Z-transform b) Obtain the in 5. a) Solve the dia and $x(1) = 2$ b) State and pro-	$d\theta$ d θ by contour + cos θ he Z-transform of co of a ^k cos k θ . werse Z-transform of fference equation x(k and u(k) is unit step for	integration in the os k θ is $\frac{z(z-z)^2}{z^2-2z}$ $\frac{2z^3+z}{(z-2)^2(z-1)}$, u +2) - x(k+1) unction. f z-transform.	$\frac{\cos \theta}{\cos \theta + 1}$. Use $\sin \theta = 1$ $\sin \theta = 1$ $\sin \theta = 1$ $\sin \theta = 1$ \sin	e. e this result to action method.` u(k) where x((find)) = 1 	X
Evaluate $\int_{0}^{2\pi} \frac{1}{2}$ 4. a) Show that t Z-transform b) Obtain the in 5. a) Solve the dia and $x(1) = 2$ b) State and pro 6. Derive one-dime	$d\theta$ d θ by contour + cos θ he Z-transform of co of a ^k cos k θ . werse Z-transform of fference equation x(k and u(k) is unit step for ove shifting theorem o	integration in the ps k θ is $\frac{z(z-z)}{z^2-2z}$ $\frac{2z^3+z}{(z-2)^2(z-1)}$, where $z = -2z^3$ +2) - x(k+1) unction. f z-transform. In governing transform.	$\frac{\cos \theta}{\cos \theta + 1}$. Use $\sin \theta = 1$ Use $\sin \theta = 1$ Use $\sin \theta = 1$ Use $\sin \theta =$	e. this result to ction method.` u(k) where x((n of string and	 find)) = 1 solve 	
Evaluate $\int_{0}^{2\pi} \frac{1}{2}$ 4. a) Show that t Z-transform b) Obtain the in 5. a) Solve the dia and $x(1) = 2$ b) State and pro- it completely.	$\frac{d\theta}{d\theta}$ d θ by contour the Z-transform of co of a ^k cos k θ . werse Z-transform of fference equation x(k and u(k) is unit step for ove shifting theorem o ensional wave equation	integration in the os k θ is $\frac{z(z-z)^2}{z^2-2z}$ $\frac{2z^3+z}{(z-2)^2(z-1)}$, u +2) - x(k+1) unction. f z-transform. n governing transform	$\frac{\cos \theta}{\cos \theta + 1}$. Use $\sin \theta + 1$. Use $\sin \theta + 1$ = $0.25x(k) =$	e. e this result to action method.` u(k) where x(() n of string and) find)) = 1 solve 	
 Evaluate ∫₀^{2π}/₂ 4. a) Show that t Z-transform b) Obtain the in 5. a) Solve the dia and x(1) = 2 b) State and prof. 6. Derive one-dimensit completely. 	$\frac{d\theta}{d\theta}$ d θ by contour $\frac{d\theta}{d\theta}$ by contour he Z-transform of co of $a^k \cos k\theta$. werse Z-transform of fference equation x(k and u(k) is unit step for ove shifting theorem o ensional wave equation	integration in the os k θ is $\frac{z(z-z)}{z^2-2z}$ $\frac{2z^3+z}{(z-2)^2(z-1)}$, u +2) - x(k+1) unction. f z-transform. n governing transform.	$\frac{\cos\theta}{\cos\theta+1}$. Use $\sin\theta$ sing partial fra + 0.25x(k) = sverse vibration	e. e this result to action method.` u(k) where x((n of string and	 find)) = 1 solve 	· · · · · · · · · · · · · · · · · · ·

- 7. Solve the one dimensional heat equation $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$ under the conditions:
 - a) u is not infinite as $t \rightarrow \infty$
 - b) $\frac{\partial u}{\partial x} = 0$ for x = 0 and x = l and
 - c) $u(x,0) = lx x^2$ for t = 0 between x = 0 and x = l

OR

The diameter of a semi circular plate of radius a is kept at 0°C and temperature at the semi circular boundary is T°C. Show that the steady temperature in the plate is given

by u(r,
$$\theta$$
) $\frac{4T}{\pi} \sum_{n=1}^{\infty} \frac{1}{2n-1} \left(\frac{r}{a}\right)^{2n-1} \sin(2n-1)\theta$

& a) Find the Fourier cosine integral representation of the function $f(x) = e^{-kx}$ (x > 0, k > 0) and hence show that

$$\int_{0}^{\infty} \frac{\cos \omega x}{k^{2} + \omega^{2}} d\omega = \frac{\pi}{2k} e^{-kx} \quad (x > 0, K > 0)$$

b) Obtain Fourier sine transform of e^{-x} , (x > 0) and hence evaluate $\int_{0}^{\infty} \frac{x^2}{(1+x^2)^2} dx$.

Exam.	Regular / Back		
Level	BE	Full Marks	80
Programme	BEL, BEX, BCT	Pass Marks	32
Year / Part	П/П	Time	3 hrs.

2067 Mangsir

Subject: - Applied Mathematics

✓ Candidates are required to give their answers in their own words as far as practicable.

Attempt any <u>Six</u> questions.

✓ <u>All</u> questions carry equal marks.

Assume suitable data if necessary.

- 1. a) State Cauchy Riemann equations in polar form. Show that $f(z) = \sin z$ is analytic in the entire z-plane.
 - b) State and prove Cauchy's integral formula.
- 2. a) State Laurent series. Find Taylor series of $f(z) = \cos z$ about $z = \frac{\pi}{4}$.

b) Define pole of order m. Find the residue of $f(z) = \frac{Z^2 e^z}{(Z-2)^3}$ at its pole.

3. a) Determine the Z-transform of

, i) t²e^{-at}

- ii) e^{-at} coswt
- b) State initial value theorem for Z-transform. If Z-transform of a function is given by

$$X(z) = \frac{(1 - e^{-t})z^{-1}}{(1 - z^{-1})(1 - e^{-t}z^{-1})}, \text{ determine } x(0), x(1) \text{ and } x(2).$$

- 4. a) Find inverse Z- transform of
 - i) $x(z) = \frac{z+2}{z^2 5z + 6}$ (by partial fraction method)
 - ii) $x(z) = \frac{z+2}{z^2 + 7z + 10}$ (by inversion integral method)
 - b) Solve the difference equation: x(k+2) 4x(k+1) + 4x(k) = 0 Where x(o) = 1 and x(1) = 0.
- 5. Derive one dimensional wave equation and obtain its solution.
- 6. Solve: $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$ subject to the conditions, $u(o,y) = u(\ell,y) = u(x,o) = 0$, and $u(x,a) = \sin\left(\frac{n\pi x}{\ell}\right)$.
- 7. Define convolution for Fourier transform. Verify convolution theorem for $f(x) = g(x) = e^{-x^2}$.
- 8. Maximize: $z = x_1 + 3x_2$ subject to

 $x_1 + 2x_2 \le 10, x_1 \le 5$, and $x_2 \le 4; x_1, x_2 \ge 0$

by using simplex method.